

The XMM-*Newton* view of the relativistic spectral features in AXJ0447-0627

R. Della Ceca¹, L. Ballo², V. Braito¹, and T. Maccacaro¹

¹ *INAF - Osservatorio Astronomico di Brera, via Brera 28, 20121 Milan, Italy*
(*rdc@brera.mi.astro.it, braito@brera.mi.astro.it, tommaso@brera.mi.astro.it*)

² *SISSA/ISAS, International School for Advanced Studies, via Beirut 4, 34014 Trieste, Italy* (*ballo@sissa.it*)

ABSTRACT

The XMM-*Newton* observation of the optically Type 1 AGN AXJ0447-0627 ($z = 0.214$) unambiguously reveals a complex, bright and prominent set of lines in the 4 – 8 keV rest frame energy range. Although, from a phenomenological point of view, the observed properties can be described by a simple power law model plus 5 narrow Gaussian lines (at rest frame energies of ~ 4.49 , ~ 5.55 , ~ 6.39 , ~ 7.02 and ~ 7.85 keV), we find that a model comprising a power law ($\Gamma \sim 2.2$), a reflected relativistic continuum, a narrow Fe I $K\alpha$ line from neutral material as well as a broad Fe $K\alpha$ relativistic line from a ionized accretion disk represents a good physical description of the data. The “double horned” profile of the relativistic line implies an inclination of the accretion disk of $\sim 45^\circ$, and an origin in a narrow region of the disk, from $R_{\text{in}} \sim 19 GM/c^2$ to $R_{\text{out}} \sim 30 GM/c^2$. The narrow Fe I $K\alpha$ line from neutral material is probably produced far from the central black hole, most likely in the putative molecular torus. Although some of these properties have been already found in other Type 1 AGN and discussed in the literature, at odd with the objects reported so far we measure high equivalent widths (EWs) of the observed lines: ~ 1.4 keV for the “double horned” relativistic line and ~ 0.4 keV for the narrow line.

Subject headings: galaxies: active – galaxies: individual (AXJ0447-0627) – galaxies: Seyfert – X-rays: galaxies

1. Introduction

Recent XMM-*Newton* and *Chandra* observations of a number of Active Galactic Nuclei have discovered complex spectral features (broad and/or narrow) red-ward of the well known

fluorescent Fe emission line(s) at 6.4–7 keV (Turner et al. 2002; Guainazzi 2003; Yaqoob et al. 2003; Turner et al. 2004; Porquet et al. 2004; Bianchi et al. 2004; McKernan & Yaqoob 2004; Gallo et al. 2004). These features are probably produced in (or very close to) the supposed accretion disk and, as such, their study provides primary information about the dynamics and physical processes which are taking place in the innermost part of the AGN (see Fabian, Iwasawa, Reynolds, & Young 2000 and Reynolds & Nowak 2003 for a review). Among the models invoked to explain these features there are: *i*) localized hot spot on the accretion disk surface due to illumination by local flares (e.g. Dovčiak et al. 2004); *ii*) reprocessed emission from narrow annuli on the surface of the accretion disk (Gallo et al. 2004); *iii*) inflow or outflow of material (e.g. the ejected blob model proposed by Turner, Kraemer, & Reeves 2004 to explain the properties of the narrow lines observed in MKN 766); and *iv*) destruction (spallation) by energetic protons on the accretion disk surface of Fe into lower Z elements (mainly Cr and Mn), implying an enhancement of the line emission expected from elements of lower abundance (Skibo 1997).

In this paper we use XMM-*Newton* data to discuss the observed line properties of AXJ0447-0627, a broad line AGN at $z = 0.214$, discovered during the optical identification process of the X-ray sources of the *ASCA* (2 – 10 keV) Hard Serendipitous Survey (Cagnoni, Della Ceca, & Maccacaro 1998; Della Ceca et al. 1999). We assume $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\lambda = 0.7$ and $\Omega_M = 0.3$.

2. Observations and data reduction

2.1. X-ray observations

AXJ0447-0627 was detected with a $S/N = 5.40$ in the *ASCA* field pointed at NGC 1667 (*ASCA* Sequence ID= 71032000) at the nominal *ASCA* position of RA = 04:47:48.6, Dec = –06:27:50.8 (J2000.0). The net *ASCA* GIS2 counts from the source are 54 ± 10 (2 – 10 keV), corresponding to a $f_{2-10 \text{ keV}} \sim 3 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (assuming a power-law photon index $\Gamma = 1.7$). The source attracted our attention because of its position in the hardness ratio diagram (cf. Della Ceca et al. 1999), indicative of a hard, presumably obscured, X-ray source ($HR1 = -0.40 \pm 0.15$; $HR2 = 0.32 \pm 0.17$).

We thus observed AXJ0447-0627 with XMM-*Newton* on Sept 8, 2002 for a total of about 29 ks. The three EPIC cameras (MOS1, MOS2 and pn) were operating in full frame mode and with the thin filter applied. The XMM-*Newton* data have been cleaned and processed using the Science Analysis Software (SAS version 5.4), and analyzed using standard software packages (FTOOLS version 4.2, XSPEC version 11.2). Event files have been thus filtered

for high-background time intervals and only events corresponding to patterns 0 – 12 (MOS 1&2) and 0 – 4 (pn) have been used (see Ehle et al. 2001); the net exposure times at the source position after data cleaning are ~ 21.4 ks (MOS1, MOS2) and ~ 17.5 ks (pn).

The XMM-*Newton* MOS1, MOS2 and pn images in the 0.5 – 10 keV energy range reveal a high signal-to-noise ratio ($S/N \sim 30$ and ~ 50 in the MOS and the pn, respectively) point-like source within the *ASCA* 90% positional error circle ($\sim 2'$ radius) of AXJ0447-0627. This is the only detected and visible X-ray source in the *ASCA* error circle; the X-ray position derived using the XMM-*Newton* data is RA = 04:47:48.62, Dec = $-06:28:12$ ($\sim 21''$ away from the nominal *ASCA* position).

Source counts were extracted from a circular region of radius $22.5''$ for the MOS and $17.5''$ for the pn (this smaller radius for the pn is due to the proximity of a CCDs gap). Background counts were extracted from a nearby source-free circular region of $\sim 42'' - 50''$ radius. The net count rates (0.5 – 10 keV energy range) are 0.051 ± 0.002 counts s^{-1} , 0.054 ± 0.002 counts s^{-1} and 0.171 ± 0.004 counts s^{-1} for MOS1, MOS2 and pn, respectively; the source counts represents about 85% of the total counts in the source extraction region. No statistically significant source variability has been detected during the XMM-*Newton* observation. To improve statistics, the MOS1 and MOS2 data have been combined together, and the MOS and pn spectra have been fitted simultaneously, keeping the relative normalization free. Source counts were binned so as to have at least 20 counts in each energy bin. We have also generated our own spectral response matrices at the source position using the SAS tasks *arfgen* and *rmfgen*. All the models discussed here have been filtered through the Galactic absorption column density along the line of sight ($N_{H, Gal} = 5.6 \times 10^{20} \text{ cm}^{-2}$; Dickey & Lockman 1990). Unless otherwise stated, fit parameters are quoted in the rest-frame of AXJ0447-0627 ($z = 0.214$, see below), while the figures and the EWs are in the observer frame.

2.2. Source identification and optical spectroscopy

A bright (RA = 04:47:48.5, Dec = $-06:28:13$; APM red magnitude = 17.7) optical source lies about $2''$ from the X-ray position derived using the XMM-*Newton* data. This object was observed spectroscopically at the TNG on October 5th, 2002. The optical spectrum (not reported here) covers the wavelength range $\sim 3500 - 8000 \text{ \AA}$ (dispersion of $2.8 \text{ \AA/pixel}$) and clearly shows broad (FWHM $> 6000 \text{ km s}^{-1}$) MgII, H β and H α lines as well as narrow (FWHM $< 1000 \text{ km s}^{-1}$) [OIII]4959,5007 $\text{\AA}$ lines. The optical line properties and position allow us to classify AXJ0447-0627 as a classical broad line AGN at $z = 0.214 \pm 0.001$. AXJ0447-0627 also belongs to the *ASCA* Medium Sensitivity Survey, so an independent

confirmation of the redshift and of the optical spectral classification comes from the work presented in Akiyama et al. (2003).

3. X-ray spectral analysis

A single absorbed power law (PL) model is not a good description of the overall (0.5 – 10 keV) spectrum of AXJ0447-0627 ($\Gamma = 2.18 \pm 0.05$; N_{H} consistent with zero; $\chi^2/\text{d.o.f} = 280.3/250$), since a very large discrepancy is present above 4 keV.

In Fig. 1 we show the ratio between the best fit PL model (obtained considering only the line-free region from ~ 0.8 to ~ 3 keV) and the data in the 2 – 10 keV energy range. The residuals show the presence of a possible “line-like” feature at $E \sim 3.5$ keV (observer frame), and of a complex structure (several different lines?) in the energy range between ~ 4.5 and ~ 7 keV (observer frame). Splitting the total observation into two intervals of similar exposure times, we do not find convincing evidence of temporal variability of such structures.

We have been unable to reproduce the complex lines structure with a pure reflected continuum, either normal (PEXRAV model in XSPEC; see Magdziarz & Zdziarski 1995) or relativistic (REFSCH model; see Magdziarz & Zdziarski 1995; Fabian, Rees, Stella, & White 1989). A pure reflected continuum with associated emission lines from Ca, Cr, Fe and Ni (see e.g. the modeling of the Seyfert 2 galaxy NGC 6552 by Reynolds et al. 1994) is also unable to reproduce the observed structure since the relative abundances of the above elements are different from the expected ones. We also note that a pure reflected continuum, more typical of optical Type 2 AGNs, is at odd with the optical spectral classification of AXJ0447-0627.

To investigate the presence of line-like features and to locate their energy centroid we slid a narrow ($\sigma = 0.1$ keV) Gaussian template across the data between 4 and 8 keV (rest frame), looking for an improvement to the fit with respect to the simple PL model. The results are shown in the inset of Fig. 1, where we report the change in fit statistics ($\Delta\chi^2$) as a function of the centroid energy position of the narrow Gaussian line. This analysis points out the presence of a number of possible narrow lines, with rest frame energy centroids at about 4.5, 5.6, 6.4, 7.0 and 7.9 keV (in the last case, the line centroid is not well constrained). Note that in the resulting fit statistics a $\Delta\chi^2$ greater than 10.30 (5.25) represents a feature that is significant at more than 99% (90%) confidence level, so the lines at energies ~ 4.5 keV and at ~ 7.9 keV are of lower statistical significance with respect to those at ~ 5.6 keV, ~ 6.4 keV and ~ 7.0 keV. Following these indications, we tried to reproduce the observed spectrum with a power law and five narrow Gaussian lines. The best fit spectral parameters

are reported in Table 1 while the ratio between the data and this possible best fit model is shown in Fig. 2 (panel a).

The line at $6.39^{+0.07}_{-0.06}$ keV ($\text{EW} \sim 700$ eV) is positionally consistent with the Fe I $K\alpha$ emission line, while the line at $7.02^{+0.29}_{-0.12}$ keV ($\text{EW} \sim 600$ eV) is positionally consistent both with the Fe I $K\beta$ emission line (rest frame energy $E = 7.058$ keV) or with the Fe XXVI $\text{Ly}\alpha$ emission line (rest frame energy $E = 6.96$ keV). However the association with the Fe I $K\beta$ line is unlikely since the measured flux from this line should be at a fixed ratio (~ 0.11) with the Fe I $K\alpha$ emission flux, clearly in disagreement with the measured EWs. The association with Fe XXVI $\text{Ly}\alpha$ seems to be more plausible given that this line could be very prominent (and sometimes with an EW comparable with the narrow Fe I $K\alpha$ line) in Type 1 AGN (see e.g. the case of the Seyfert 1 galaxy NGC 7314 discussed in Yaqoob et al. 2003). The line at $7.85^{+0.73}_{-0.76}$ keV is positionally consistent with the Ni $K\alpha$, while for the remaining two lines ($E = 4.49^{+0.13}_{-0.17}$ keV and $5.55^{+0.06}_{-0.05}$ keV) there are no clear associations with well known and expected elements. The strongest expected lines in the spallation model are the Cr $K\alpha$ at 5.4 keV and the Mn $K\alpha$ at 5.9 keV (Skibo 1997). Both these lines are ruled out by the mismatch with the measured energy lines centroid. So, unless an energy shift occurs (but we do not observe any energy shift for the Fe I $K\alpha$ line), the spallation model is an unlikely explanation of the XMM-*Newton* data. We have also evaluated the upper limits for Fe XXV (f) at $E \sim 6.64$ keV and for Fe XXV (r) at $E \sim 6.70$ keV since the strenght of these lines, when combined with the strenght of other ionized Fe lines, can constrain emission models (cfr. Yaqoob & Padmanabhan 2004 and reference therein). These two lines are not required by the current data set and the 90% upper limit on their EW is ~ 400 eV.

The complex structure detected in the spectrum of AXJ0447-0627 could suggest a profile of a Fe line produced by an accretion disk. We explored this interesting possibility using the DISKLINE model (Fabian, Rees, Stella, & White 1989), which assumes a non-rotating Schwarzschild black hole¹. The relativistic effects have been introduced also in the description of the reflected continuum, replacing the simple power law model with the XSPEC model REFSCH².

We started the analysis using a model composed of the REFSCH model plus a Fe I $K\alpha$

¹We have also tried the LAOR model (Laor 1991), in which the black hole is maximally rotating, but because of the limited source count statistics we could not discriminate between the DISKLINE and LAOR model. Since the black hole spin parameter is clearly an over-parameterization of the present data set we report here only the results obtained applying the DISKLINE model. We note however that similar results have been obtained using the LAOR model.

²This model is the sum of an e-folded power law primary spectrum plus its reflected component from a ionized relativistic accretion disk.

disk line fixing its energy position to 6.4 keV rest-frame; since in AXJ0447-0627 the observed lines seems to have a significantly larger EW than usual we have also added the corresponding Fe I $K\beta$ disk line emission fixing the Fe I $K\beta$ /Fe I $K\alpha$ ratio to that expected from the theory (~ 0.11). The best fit spectral parameters are reported in Table 2 while the ratio between the data and the best fit model is shown in Fig. 2 (panel b). Although the overall fit is statistically acceptable ($\chi^2_\nu = 1.00$), the ratio in Fig. 2 (panel b) shows a “line-like” residual at an observed energy of ~ 6 keV (~ 7.3 keV rest frame) that we were unable to reduce. We tried to consider also disk lines associated to Ni $K\alpha$ at ~ 7.5 keV or to Fe XXVI $\text{Ly}\alpha$ at ~ 6.96 keV emission, and/or allowed the abundance of Fe I $K\beta$, Ni $K\alpha$ and Fe XXVI $\text{Ly}\alpha$ to be a free parameter of the fit, but we were unable to take into account such “line-like” structure.

Thus we tried the same model first used by Weaver et al. (1997) to describe the spectral properties of MCG-5-23-16: a narrow Fe I $K\alpha$ component (E_{line} fixed at 6.4 keV) plus a broad relativistic line component (DISKLINE model) along with the underlying continuum (REFSCH model). The best fit spectral parameters are reported in Table 2, the ratio between the data and the best fit model is shown in Fig. 2 (panel c), while the folded model and the model itself are reported in Fig. 3. The observed line position ($E = 6.61 \pm 0.11$ keV) of the broad relativistic line is inconsistent with the Fe I $K\alpha$ from neutral material but strongly suggests that it is due to Fe $K\alpha$ emission from ionized He-like material. Overall this modeling provide a good description of the broad band spectral properties of AXJ0447-0627. We have also tried to add a relativistic emission line from Fe XXVI $\text{Ly}\alpha$ to the best fit model reported in Table 2; such line can be accommodated within the present data set (with an EW of ~ 15 eV) but it is not statistically required. The observed flux and the intrinsic luminosity in the 0.5 – 10 keV energy range are $(6.5 \pm 0.4) \times 10^{-13}$ erg cm $^{-2}$ s $^{-1}$ and $(8.9 \pm 0.5) \times 10^{43}$ erg s $^{-1}$, the errors reflecting the uncertainties on the best fit model. We note that the 2 – 10 keV flux measured with XMM-Newton ($\sim 3.6 \times 10^{-13}$ erg cm $^{-2}$ s $^{-1}$) is in very good agreement with the 2 – 10 keV flux measured with ASCA.

4. Discussion and Conclusions

Using XMM-Newton data we have revealed that the optically Type 1 AGN AXJ0447-0627 at $z = 0.214$ is characterized by a complex of bright and prominent set of lines in the 4.5 – 8.0 keV energy range (rest frame).

We have shown that these lines can be reasonably well reproduced by a physical model comprising a power law, a reflected relativistic continuum, a narrow Fe I $K\alpha$ line from neutral material and a Fe He-like $K\alpha$ relativistic line from a ionized accretion disk. Although not

well constrained, the best fit ionization parameter (ξ in Table 2) is consistent with the Fe ionization state, as deduced from the best fit line position (cf. Matt, Fabian, & Ross 1993). A similar modeling of the Fe line properties (a narrow plus a broad relativistic component) has been found to describe, for example, the spectral properties of the Seyfert 1.9 MCG-5-23-16 (Weaver et al. 1997), of the radio-quiet quasar MKN 205 (Reeves et al. 2001) and of the Seyfert 1 NGC 3516 (Turner et al. 2002); the presence of the Fe relativistic line from a high ionized accretion disk has been unambiguously reported in the case of MKN 205. As already discussed by the above authors, the most likely origin of the narrow 6.4 keV component is from neutral matter distant from the black hole, e.g. the putative molecular torus.

The resulting best fit of the relativistic “double horned” profile implies an inclination of the accretion disk of $\sim 45^\circ$, as well as that the observed lines should be produced in a narrow region of the disk from $R_{\text{in}} \sim 19 \text{ } GM/c^2$ to $R_{\text{out}} \sim 30 \text{ } GM/c^2$. As in the case of e.g. ESO 198-G24 (Guainazzi 2003) and NAB 0205+024 (Gallo et al. 2004), a few alternative possibilities can be conjectured to explain why the inner radius is larger than the last stable orbit. The first possibility is that the disk is highly ionized in the inner part, so most of the Fe is completely stripped off and the production of the Fe lines is suppressed. Second, the accretion disk in AXJ0447-0627 could be truncated at $10 - 15 \text{ } GM/c^2$ (see Müller & Camenzind 2004). Finally the relativistic line can be produced by a localized hot spot on the accretion disk surface (e.g. Dovčiak et al. 2004).

All the properties discussed above have been also observed in other AGNs and seem to be in agreement with the expectation from accretion disk theory. However there is an observed property that is very unusual in AXJ0447-0627 and makes this object unique: the very large EW observed, which is at least a factor 5 greater than that usually measured in other Type 1 AGNs (see e.g. Yaqoob & Padmanabhan 2004 and reference therein) or expected from an accretion disk around a Schwarzschild black hole (Matt et al. 1992; Matt, Fabian, & Ross 1993). According to the modelling reported in Martocchia & Matt (1996) and Miniutti & Fabian (2004), a high EW could be explained if the primary X-ray source (illuminating both the observer and the accretion disk) is located very close to a central and maximally rotating Kerr black hole. However such a combination should also imply a very high value of the reflection parameter R and a reflection dominated source, probably in disagreement with the best fit found here.

A way to solve part of these problems is to assume that the emission features appear much stronger than normal because the continuum is strongly absorbed. We have tested this possibility by adding a partial covering absorption model in front to the underlying continuum. Unless the primary AGN emission is heavily absorbed ($N_{\text{H}} \sim 10^{25} \text{ cm}^{-2}$) and therefore with signatures falling outside the XMM-*Newton* bandpass (but this is clearly at

odd with the optical spectral classification of AXJ0447-0627), the best fit absorbing N_{H} and covering fraction ($\sim 9 \times 10^{21} \text{ cm}^{-2}$ and ~ 0.2 , respectively) imply that absorption effects can not take into account the strong emission features observed.

This said, and with the caveat that the model proposed here could be not fully appropriate (e.g. we have already pointed out that a phenomenological description of the data can be also obtained by a simple power law model plus 5 narrow Gaussian lines at rest frame energies of $4.49_{-0.17}^{+0.13} \text{ keV}$, $5.55_{-0.05}^{+0.06} \text{ keV}$, $6.39_{-0.06}^{+0.07} \text{ keV}$, $7.02_{-0.12}^{+0.29} \text{ keV}$ and $7.85_{-0.76}^{+0.73} \text{ keV}$), we would like to note that the emission lines in AXJ0447-0627 have a total EW of $\sim 2 \text{ keV}$. This is an observational result and, as such, is model independent. These lines deserve further attention and a deeper investigation since any model proposed to describe the X-ray spectral properties of AXJ0447-0627 should be able to explain their large EW.

Finally we would like to note that AXJ0447-0627 was selected as a target for an XMM-*Newton* observation because of its observed *ASCA* hardness ratios, indicative of a hard, presumably obscured, X-ray source (cf. Della Ceca et al. 1999). The XMM-*Newton* observations reported here have shown that its hard X-ray colors are due to the strong line complex in the observed $4.5 - 8.0 \text{ keV}$ energy range rather than to absorption effects. There have been many claims in recent years about a substantial fraction ($\sim 10\%$) of X-ray absorbed optically classified Type 1 AGN (with strong implications for AGN unification models and synthesis of the cosmic X-ray background) based mainly on poor quality X-ray data (e.g. hardness ratios; see Willott et al. 2004). The result presented here indicates that a number of these sources, *thought* to be X-ray absorbed Type 1 AGNs on the basis of their hardness ratios, could instead be X-ray un-absorbed AGNs with substantial and complex X-ray line emission (see also Maccacaro et al. 2004).

Based on observations obtained with XMM-*Newton* (an ESA science mission with instruments and contributions directly funded by ESA Member States and the USA, NASA) and with the Telescopio Nazionale Galileo (operated in the island of La Palma by the Centro Galileo Galilei of the INAF in the Spanish Observatorio del Roque de los Muchacos of the Instituto de Astrofísica de Canarias). We would like to thank the anonymous referee for her/his useful comments that have contributed improving this paper. We thank M. Cappi, M. Dadina, A. Caccianiga, L. Maraschi, G. Matt and P. Severgnini for useful discussions. This work has received partial financial support from the Italian Space Agency (ASI grant I/R/062/02) and from the MIUR (Cofin-03-02-23).

REFERENCES

- Akiyama, M., Ueda, Y., Ohta, K., Takahashi, T., & Yamada, T. 2003, *ApJS*, 148, 275
- Bianchi, S., Matt, G., Balestra, I., Guainazzi, M., & Perola, G. C. 2004, *A&A*, 422, 65
- Cagnoni, I., Della Ceca, R., & Maccacaro, T. 1998, *ApJ*, 493, 54
- Della Ceca, R., Castelli, G., Braitto, V., Cagnoni, I., & Maccacaro, T. 1999, *ApJ*, 524, 674
- Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
- Dovčiak, M., Bianchi, S., Guainazzi, M., Karas, V., & Matt, G. 2004, *MNRAS*, 350, 745
- Ehle, M. et al. 2001, *XMM-Newton Users' Handbook*
(http://xmm.vilspa.esa.es/external/xmm_user_support/documentation/uhb_2.0/index.html)
- Fabian, A. C., Rees, M. J., Stella, L., & White, N. E. 1989, *MNRAS*, 238, 729
- Fabian, A. C., Iwasawa, K., Reynolds, C. S., & Young, A. J. 2000, *PASP*, 112, 1145
- Gallo, L. C., Boller, T., Brandt, W. N., Fabian, A. C., & Vaughan, S. 2004, *MNRAS*, 355, 330
- Guainazzi, M. 2003, *A&A*, 401, 903
- Laor, A. 1991, *ApJ*, 376, 90
- Maccacaro, T., Braitto, V., Della Ceca, R., Severgnini, P., & Caccianiga, A. 2004, *ApJ*, 617, L33
- Magdziarz, P. & Zdziarski, A. A. 1995, *MNRAS*, 273, 837
- Martocchia, A., & Matt, G. 1996, *MNRAS*, 282, L53
- Matt, G., Perola, G. C., Piro, L., & Stella, L. 1992, *A&A*, 263, 453
- Matt, G., Fabian, A. C., & Ross, R. R. 1993, *MNRAS*, 262, 179
- McKernan, B. & Yaqoob, T. 2004, *ApJ*, 608, 157
- Miniutti, G., & Fabian, A. C. 2004, *MNRAS*, 349, 1435
- Müller, A. & Camenzind, M. 2004, *A&A*, 413, 861

- Porquet, D., Reeves, J. N., Uttley, P., & Turner, T. J. 2004, *A&A*, 427, 101
- Reeves, J. N., Turner, M. J. L., Pounds, K. A., O’Brien, P. T., Boller, T., Ferrando, P., Kendziorra, E., & Vercellone, S. 2001, *A&A*, 365, L134
- Reynolds, C. S., Fabian, A. C., Makishima, K., Fukazawa, Y., & Tamura, T. 1994, *MNRAS*, 268, L55
- Reynolds, C. S. & Nowak, M. A. 2003, *Phys. Rep.*, 377, 389
- Skibo, J. G. 1997, *ApJ*, 478, 522
- Turner, T. J., et al. 2002, *ApJ*, 574, L123
- Turner, T. J., Kraemer, S. B., & Reeves, J. N. 2004, *ApJ*, 603, 62
- Turner, T. J., Kraemer, S. B., George, I. M., Reeves, J. N., & Bottorff, M. C. 2004, *ApJ*, 618, 155.
- Weaver, K. A., Yaqoob, T., Mushotzky, R. F., Nousek, J., Hayashi, I., & Koyama, K. 1997, *ApJ*, 474, 675
- Willott, C. J., et al. 2004, *ApJ*, 610, 140
- Yaqoob, T., George, I. M., Kallman, T. R., Padmanabhan, U., Weaver, K. A., & Turner, T. J. 2003, *ApJ*, 596, 85
- Yaqoob, T., & Padmanabhan, U. 2004, *ApJ*, 604, 63

Table 1: Results of the spectral analysis (0.5 – 10.0 keV in the observed frame) - Power law plus 5 narrow Gaussian lines.

Power Law		Lines			$\chi^2/\text{d.o.f.}$
Γ	Norm	$E_{\text{r.f.}}$ (keV)	Norm	EW (eV)	
(1)	(2)	(3)	(4)	(5)	(6)
$2.24^{+0.06}_{-0.05}$	$2.38^{+0.13}_{-0.06}$	$4.49^{+0.13}_{-0.17}$	$0.92^{+0.71}_u$	93^{+72}_u	226.3/240
		$5.55^{+0.06}_{-0.05}$	$2.68^{+0.94}_{-1.01}$	436^{+152}_{-164}	
		$6.39^{+0.07}_{-0.06}$	$3.13^{+0.82}_{-1.26}$	700^{+182}_{-290}	
		$7.02^{+0.29}_{-0.12}$	$2.18^{+0.91}_{-1.08}$	602^{+251}_{-299}	
		$7.85^{+0.73}_{-0.76}$	$0.74^{+0.94}_u$	263^{+331}_u	

Columns are as follow:

Column 1: power law photon index; Column 2: normalization of the power law in units of 10^{-4} photons keV $^{-1}$ cm $^{-2}$ s $^{-1}$ @ 1 keV; Column 3: rest-frame energy centroid of the narrow Gaussian line; Column 4: normalization in units of 10^{-6} photons keV $^{-1}$ cm $^{-2}$ s $^{-1}$ in the line; Column 5: equivalent width of the line; Column 6: χ^2 and number of degrees of freedom.

NOTE: Errors are quoted at the 90% confidence level for 1 parameter of interest ($\Delta\chi^2 = 2.71$);

u : unconstrained parameter.

Table 2: Results of the spectral analysis ($0.5 - 10.0$ keV in the observed frame) - REFSCHE model + Fe $K\alpha$ relativistic line + Fe (relativistic or Gaussian) line.

REFSCH ^(a)							DISKLINE			
							DISKLINE or GAUSSIAN			χ^2 [d.o.f.]
Γ	i	R_{in}	R_{out}	ξ	R	Norm	$E_{\text{r.f.}}$ (keV)	Norm	EW ^(b) (keV)	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
$2.26^{+0.04}_{-0.07}$	$25.0^{+4.4}_{-3.0} (c)$	$16.2^u_{-3.9} (d)$	$25.2^{+13.5}_u (d)$	$1f$	$0.44 - 2.00$	1.55 ± 0.05	$6.4f$	$5.3^{+1.6}_{-1.2} (c)$	$1.3^{+0.4}_{-0.3}$	235.94
<i>relativistic Fe I $K\beta$:</i>							$7.06f$	$0.6^{+0.2}_{-0.1} (c)$	$0.18^{+0.05}_{-0.04}$	[246]
$2.21^{+0.09}_{-0.15}$	$45.1^{+6.8}_{-7.4} (c)$	$19.5^u_{-6.0} (d)$	$29.5^{+102.1}_u$	$576.4^{+956.6}_u (c)$	$1.00^{+1.00}_{-0.95}$	$1.53^{+0.06}_{-0.09}$	$6.61^{+0.11}_{-0.12} (c)$	$6.2^{+2.1}_{-2.5}$	$1.4^{+0.7}_{-0.4}$	223.2
<i>narrow Fe I $K\alpha$:</i>							$6.4f$	$1.7^{+1.1}_{-1.0}$	$0.4^{+0.2}_{-0.2}$	[243]

Columns are as follow:

Column 1: power law photon index; Column 2: inclination angle (degrees); Column 3: inner disk radius in units of GM/c^2 ; Column 4: outer disk radius in units of GM/c^2 ; Column 5: disk ionization parameter in units of erg cm s^{-1} ; Column 6: reflection scaling factor; Column 7: photon flux @ 1 keV of the cutoff broken power law only (no reflection) in the observed frame in units of $10^{-4} \text{ photons keV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$; Column 8: rest-frame energy centroid of the line; Column 9: normalization in units of $10^{-6} \text{ photons cm}^{-2} \text{ s}^{-1}$ in the line; Column 10: equivalent width of the line; Column 11: χ^2 and number of degrees of freedom.

NOTE: Errors are quoted at the 90% confidence level for 1 parameter of interest ($\Delta\chi^2 = 2.71$). During the fit, the disk parameters of the different components were tied together; f : fixed parameter; u : unconstrained parameter.

^(a) The following parameters have been fixed during all the fits: cutoff energy $E_c = 100$ keV; disk temperature $T = 3 \times 10^4$ K; power law index for reflection emissivity $\beta = -3$.

^(b) Equivalent widths are computed with respect to the REFSCHE underlying continuum.

^(c) Errors were calculated fixing at the best fit values R_{in} and R_{out} (inclination of the accretion disk and relativistic line energy); R_{in} , R_{out} and Γ (relativistic line normalization); R and Gaussian line normalization (disk ionization parameter).

^(d) Errors have been evaluated performing a fit while stepping the value of R_{in} (R_{out}) through the range $6.0 - 25.0$ ($20.0 - 200.0$).

Figure Captions

Fig. 1.— Ratio between the 2 – 10 keV MOS (filled squares) and pn (open circles) data and the best fit power law model (limited to the ~ 0.8 to ~ 3 keV energy range: $\Gamma = 2.24^{+0.21}_{-0.08}$). In the inset we report the change in fit statistic ($\Delta\chi^2$) as a function of the centroid energy position of a narrow Gaussian line model that was stepped across the data; the comparison model is the underlying power law continuum.

Fig. 2.— *Panel a*: ratio between the 2 – 10 keV MOS (filled squares) and pn (open circles) data and the best fit spectral model composed by a power law ($\Gamma = 2.24^{+0.06}_{-0.05}$) plus five Gaussian lines as detailed in Table 1. *Panel b*: ratio between the data and the best fit spectral model composed by the REFSCHE model (an e-folded power law primary spectrum plus its reflected component from a ionized relativistic accretion disk) plus relativistic Fe I $K\alpha$ and Fe I $K\beta$ lines as detailed in Table 2. *Panel c*: ratio between the data and the best fit spectral model composed by the REFSCHE model plus a narrow Fe I $K\alpha$ line from neutral material and a broad Fe relativistic line from a ionized accretion disk, as detailed in Table 2.

Fig. 3.— MOS (filled squares) and pn (open circles) folded spectra fitted with the REFSCHE model plus a narrow Fe I $K\alpha$ line from neutral material and a broad Fe relativistic line from a ionized accretion disk. In the inset we show the best fit model as detailed in Table 2.





